

# Record-breaking earthquake intervals in a global catalog and an aftershock sequence

Yoder, M.R.<sup>(1)</sup>, Turcotte, D.L.<sup>(2)</sup> and Rundle, J.B.<sup>(1,2,3)</sup>

- (1) Dept. of Physics, University of California Davis, Davis, CA, USA  
email mark.yoder@gmail.com
- (2) Dept. of Geology, University of California Davis, Davis, CA, USA
- (3) Santa Fe Institute, Santa Fe, NM, USA

For the purposes of this study, an interval is the elapsed time between two earthquakes in a designated region; the minimum magnitude for the earthquakes is prescribed. A record-breaking interval is one that is longer (or shorter) than preceding intervals; a starting time must be specified. We consider global earthquakes with magnitudes greater than 5.5 and show that the record-breaking intervals are well approximated by a Poissonian (random) theory. We also consider the aftershocks of the 2004 Parkfield earthquake and show that the record-breaking intervals are approximated by very different statistics. In both cases, we calculate the number of record-breaking intervals  $n_{rb}$  and the record-breaking interval durations  $t_{rb}$  as a function of "natural time", the number of elapsed events. We also calculate the ratio of record-breaking long intervals to record-breaking short intervals as a function of time,  $r(t)$ , which is suggested to be sensitive to trends in noisy time series data. Our data indicate a possible precursory signal to large earthquakes that is consistent with accelerated moment release (AMR) theory.